

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Trends in life satisfaction in European and North-American adolescents from 2002 to 2010 in over 30 countries

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1521082> since 2016-06-26T21:48:13Z

Published version:

DOI:10.1093/eurpub/ckv014

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

This is the author's final version of the contribution published as:

Cavallo F, Dalmaso P, Ottová-Jordan V, Brooks F, Mazur J, Välimaa R, Gobina I, Gaspar de Matos M, Raven-Sieberer U; Positive Health Focus Group.

Trends in life satisfaction in European and North-American adolescents from 2002 to 2010 in over 30 countries.

Eur J Public Health. 2015 Apr;25 Suppl 2:80-2. doi: 10.1093/eurpub/ckv014.

The publisher's version is available at:

[http://eurpub.oxfordjournals.org/content/25/suppl_2/80.long]

When citing, please refer to the published version.

Trends in life satisfaction in European and North-American adolescents from 2002 to 2010 in over 30 countries.

Franco Cavallo¹, Paola Dalmasso¹, Veronika Ottová-Jordan², Fiona Brooks³, Joanna Mazur⁴, Raili Välimaa⁵, Inese Gobina⁶, Margarida Gaspar de Matos⁷, Ulrike Raven-Sieberer² and the Positive Health Focus Group

1 Department of Public Health and Pediatrics, University of Torino, Italy

2 Department of Child and Adolescent Psychiatry, Psychotherapy, and Psychosomatics, Research Unit Child Public Health, University Medical Center Hamburg-Eppendorf, Germany

3 Head of Adolescent and Child Health Research, CRIPACC, University of Hertfordshire, UK

4 Institute of Mother and Child, Warsaw, Poland

5 Department of Health Sciences, University of Jyväskylä, Finland

6 Department of Public Health and Epidemiology, Riga Stradiņš University, Riga, Latvia

7 University of Lisbon, Portugal & CMTD/ Institute of Hygiene and Tropical Medicine

Corresponding author:

Prof. Franco Cavallo, MD, Department of Public Health and Pediatrics, University of Torino, Via Santena, 5-bis, 10126 Torino (Italy), Tel.: +390116705811, Fax: +390112365811, e-mail:

franco.cavallo@unito.it

Abstract

Background: Life Satisfaction (LS) is an indicator which is widely used for assessing the perception of a child's feeling about his life.

Methods: LS is assessed in HBSC via the Cantril ladder with 10 steps indicating the worst and best possible life. This range of values (0-10) was dichotomized into 'low' (0-5) vs. 'high' (6-10). Countries, age groups and genders were compared based on the Odds Ratio (OR) of declaring a higher LS in 2010 with respect to 2002.

Results: Analysing the difference between 2002 and 2010, 6 countries from Western Europe show decreasing LS: Austria, Canada, Switzerland, Denmark, Finland and Greenland. In contrast, a group of Eastern European Countries, i.e. Estonia, Croatia, Lithuania, Latvia, Russia and Ukraine, show a significant increase in LS. Data on gender and age differences confirm the lower rating of LS in girls and a decreasing rating with age.

Conclusion: The LS scale appears to be a tool capable of discriminating the level of wellbeing of adolescent population among countries.

Key words: life satisfaction, adolescents, HBSC, trend.

Introduction

Life satisfaction (LS), as an evaluation of an individual's quality of life is an important aspect of well-being¹ and one that is closely linked to subjective health.² Well-being in childhood is associated with social competence and good coping skills that lead to more positive outcomes in adulthood.³

LS in young people is strongly influenced by a dynamic interaction between their environment, which includes physical environment, housing quality, socio-economic condition, and the quality of their social and familial relationships. Key protective factors for a good LS include a sense of parent/family connectedness, with social support being supplied by at least one caring adult, good family communication⁴ and supportive peers, who can help to adjust to new situations and face stressful life events.⁵

Determinants of adolescent LS were not studied until early 1990s.⁶ LS is a global assessment of one's life and is thought to be relatively stable over time, compared with spontaneous feelings related to one's immediate experiences.⁷ Among adults it is inversely associated with depression, anxiety, suicide, work disability, fatal accidents and all-cause mortality.^{8, 9, 10, 11} Studies of LS have found that during adolescence, LS is strongly influenced by life experiences and relationships, particularly within the context of the family.^{12, 13}

The aim of our study was therefore to examine trends of LS from 2002 to 2010 in the countries covered by HBSC, to generate hypothesis concerning positive and negative trends observed and identifying possible 'clusters' of countries following the same pathway.

In this study, the underlying assumption is that change in adolescent LS across the last decade might be, at least partially, influenced by macro socio-economic conditions during this period.

Methods

Data from the Health Behaviour in School-aged Children (HBSC) 2002, 2006 and 2010 surveys have been used for trends analysis with the aim to explore differences and commonalities among different groups of countries.

The HBSC study has been collecting cross-sectional data on nationally representative samples of 11-, 13-, and 15-year olds since 2001/02 in more than 30 countries in Europe and North America.

Details on the general methodology of the HBSC survey have been published elsewhere.¹⁴

Among the 42 countries in the 2010 survey, only 31 (including Flemish Belgium examined independently, and Scotland, Wales and England as separate countries) participated in all three surveys and were included in the analyses.

The variable relative to adolescents' LS was represented by a ladder¹⁵ with steps going from 0, the lowest, to 10, the highest. Participants were asked to evaluate their LS using this visual analog scale by indicating the step on the ladder that corresponded to their satisfaction with their life. Data were analysed for the three surveys, in conjunction with age and gender of the adolescents. For the analyses, the range of possible values (0-10) was dichotomized into 'low' (0-5) vs. 'high' (6-10).

The comparison among countries and among age and gender groups was based on the computation , separately for each country, of the age and gender standardized prevalence and of the Odds Ratio of reporting a higher LS in 2010 with respect to 2002, and for the two periods separately, 2006 vs. 2002 and 2010 vs. 2006.

Data were modelled using a multivariable logistic regression where LS (dichotomized into 'low' and 'high') was the dependent variable and survey year, gender (males taken as reference) and age (11 year old school-students taken as reference) the independent ones. As the computed ORs are just relative measures of difference between one period and the other, nothing can be said about absolute differences between the two periods, or about absolute differences among countries.

A p-value for each OR was computed, presenting significance at the traditional values of 0.05, 0.01 and 0.001.

All analyses were performed using STATA v12.1 (StataCorp, College Station, TX:StataCorp LP).

Results

Observed trends for LS are presented and further discussed in terms of the relative change between one period and the other within countries, and of the observed differences among countries; absolute values were not taken into consideration, as the focus of the paper is on comparing trends within countries and among them.

Analysing the overall difference between 2002 and 2010, LS decreased for a group of 6 relatively affluent Western countries (Austria, Canada, Switzerland, Denmark, Finland and Greenland) and two belonging to the former Eastern European Countries (Hungary and Macedonia). In contrast, increasing LS was observed in a group of Eastern European Countries, i.e. Estonia, Croatia, Lithuania, Latvia, Russia and Ukraine, and in four Western European countries (Spain, Norway, Portugal and Belgium).

Analysing the two periods separately (2002-2006 and 2006-2010) revealed that for Greenland and Hungary the decreasing trend was mainly determined by the first period, while for the other decreasing countries the significant reduction occurred between 2006 and 2010. There was only a steady decrease in reported LS across all three time points in Macedonia (ORs: 0.798 and 0.867 respectively) and Switzerland (ORs: 0.924 and 0.916 respectively).

As for the 'high satisfaction countries' belonging to the Western European cluster, the exception occurred among Spanish adolescents, with a drop in ORs from 1.446 in the first period to 0.837 in the second.

In the Czech Republic, which has been classified as seeing no change overall, the opposite has occurred, with a decrease between 2002 and 2006 ($OR=0.864$), followed by an increase between 2006 and 2010 ($OR=1.298$).

In terms of gender differences, females have, with a few exceptions (Latvia, Macedonia and Ukraine), a general and significant tendency to a lower level of LS in all countries.

LS also decreases with increasing age, even if this pattern has many exceptions in the surveyed countries. The sharp decrease occurs in most countries at 15 years, with the only exception of Canada and Czech Republic, where the rating of LS remains unchanged across age. At 13 years the decrease in rating is less marked, and 6 countries (Canada, Czech Republic, Hungary, Italy, Latvia, and Ukraine) do not show significant differences with respect to their fellow mates of 11 years.

Discussion

The strength of this study lies on the quantity and quality of data, collected in comparable ways and with similar protocols in all involved countries, allowing, for the first time, to have a cross-national view of a decennial trend in LS of youth in Europe and North America.

The main limitation results from the fact that the comparison between countries is based on relative changes, not on absolute levels. A country with a sharp decline from a high value may still end up having higher LS than a country which starts with a low value and shows a sharp increase. For the aim of the study, this might not be a crucial drawback, as the focus is pointed on the analysis of changing trends within a country, and on the different trends observed in different sets of countries behaving in a similar way, with a view of the possible relation with on-going macro socio-economic conditions.

Trends in LS across Europe and North America show a quite scattered picture. In fact, increasing, decreasing and stable situations are split in a similar way (12 countries with increasing values of LS, 7 with decreasing values and 12 with stable ones).

It is interesting to note a 'Northern European' cluster of countries characterized by decreasing LS between 2002 and 2010, and an 'East European' cluster with increasing LS. Spain seems to be the only country with a decrease in LS that appears to follow the general economic crisis.

Some similarities with the trend in self-rated health¹⁶ can be pointed out as, for example, the similar trend in Hungary and Greenland (worst rating in the period 2002-2006), the marked trend of the Czech Republic towards increasing values in the period 2006-2010 and that of Spain and Denmark in the opposite direction in the same time interval.

Data on gender and age differences confirm what has already been evidenced in the recent literature, namely that girls report a lower LS than boys and that LS decreases with increasing age.¹⁷ In addition, the trend towards decreasing satisfaction with increasing age is consistent across all countries and across age.

The crucial transition towards lower LS seems to take place between 13 and 15 yrs, as if the onset of adolescence were the crucial event, rather than the physiological change taking place, especially in girls, usually at an earlier age.

As for the geographical cluster, no clear pattern seems to emerge. A more detailed analysis of their characteristics, including socio-economic data and cultural features at the national level, would be necessary to gain a deeper insight into the variability of this phenomenon.

In conclusion, the LS scale appears to be a tool that is not only capable of discriminating the level of wellbeing among countries, but also of catching the qualities of emotional well-being that are different from those captured by measuring self-rated health. Using these two indicators together is therefore likely to be of value for public health practitioners for the overall assessment of the health of the adolescent population.

Acknowledgements

The Health Behaviour in School-aged Children (HBSC) study is an international study carried out in collaboration with the WHO/EURO. The international coordinator of the study was Candace Currie, University of St. Andrews, Scotland. The data bank manager of the study was Oddrun Samdal, University of Bergen, Norway. A complete list of participating countries and researchers is available on the HBSC website (<http://www.hbsc.org>). We thank all pupils, teachers and scientific co-workers who participated in the HBSC surveys.

Funding

The data collection for each HBSC survey is funded at the national level. Financial support is provided by various government ministries, research foundations and other funding bodies.

Conflicts of interest

None declared.

Keypoints

- This is the first cross-national study of a decennial trend in Life Satisfaction of youth in Europe and North America.
- A 'Northern European' cluster of countries characterized by decreasing LS between 2002 and 2010, and an 'East European' cluster with increasing LS can be noted.
- Girls report a lower LS than boys and a decreasing LS with increasing age, across all countries and across age.
- The LS index is likely to be of value for public health practitioners for the overall assessment of the health of the adolescent population.

References

- ¹Diener E, Suh EM, Lucas RE, Smith HL. Subjective well-being: three decades of progress. *Psychol Bull* 1999; 125:276-302.
- ²Huebner ES, Suldo SM, Smith LC, McKnight CG. Life satisfaction in children and youth: empirical foundations and implications for school psychologists. *Psychol Schools* 2004; 41:81-93.
- ³Morgan A, Currie C, Due P, et al. Mental well-being in school-aged children in Europe: associations with social cohesion and socioeconomic circumstances. In: *Social cohesion for mental well-being among adolescents*. Copenhagen, WHO Regional Office for Europe, 2008:12-25.
- ⁴Levin KA, Currie C. Family structure, mother-child communication, father-child communication, and adolescent life satisfaction. A cross-sectional, multilevel analysis. *Health Education* 2010; 110:152-68.
- ⁵Schneider BH. *Friends and enemies: peer relations in childhood*. London: Arnold; 2000.
- ⁶Suldo SM, Riley KN, Shaffer EJ. Academic Correlates of Children Adolescents' Life Satisfaction. *Sch PsycholInt* 2006; 27: 567-82.
- ⁷Pavot WG, Diener E. Review of the Satisfaction with Life Scale. *Psychol Assessment* 1993; 5: 164-72.
- ⁸Fiscella K, Franks P. Does psychological distress contribute to racial and socioeconomic disparities in mortality? *Soc Sci Med* 1997; 45: 1805-9.
- ⁹Helliwell JF. Well-being and social capital: Does suicide pose a puzzle? *Soc Indic Res* 2007; 81: 455-96.

-
- ¹⁰Koivumaa-Honkanen HT, Honkanen RJ, Koskenvuo M, Viinamaki H, Kaprio J. Life dissatisfaction as a predictor of fatal injury in a 20-year follow-up. *Acta Psychiat Scand* 2002; 105: 444–50.
- ¹¹Koivumaa-Honkanen HT, Koskenvuo M, Honkanen RJ, Viinamaki H, Heikkilä K, Kaprio J. Life dissatisfaction and subsequent work disability in an 11-year follow-up. *Psychol Med* 2004; 34:221–8.
- ¹²Gohm CL, Oishi S, Darlington J, Diener E. Culture, parental conflict, parental marital status, and the subjective well-being of young adults. *J Marriage Fam* 1998; 60:319–34.
- ¹³Rask K, Astedt-Kurki P, Paavilainen E, Laippala P. Adolescent subjective well-being and family dynamics. *Scand J Caring Sci* 2003; 17: 129–38.
- ¹⁴Currie C, NicGabhainn S, Godeau E. The Health Behaviour in School-aged Children: WHO Collaborative Cross-National (HBSC) study: origins, concept, history and development 1982–2008. *Int J Public Health* 2009; 54(2):131–9.
- ¹⁵Cantril H. The pattern of human concerns. New Brunswick, NJ: Rutgers University Press, 1965.
- ¹⁶Cavallo F, Dalmaso P, Ottova V, et al. Trends in Self-rated Health in European and North-American adolescents from 2002 to 2010 (*EJPH, this issue*).
- ¹⁷Cavallo F, Zambon A, Borraccino A, Raven-Sieberer U, Torsheim T, Lemma P. Girls growing through adolescence have a higher risk of poor health. *Qual Life Res* 2006; 15: 1577–85.

Tables
Table 1 –Age and Gender Standardized Prevalence[§] and Odds Ratios for Life Satisfaction
(dichotomised in ‘low’ vs. ‘high’) for each country.

	Age and Gender			Odds Ratios					
	Standardized Rate[§]			(Adjusted for survey year, gender and age group)					
	(per 100 Children)								
	<i>2002</i>	<i>2006</i>	<i>2010</i>	<i>2010 vs</i>	<i>2006 vs</i>	<i>2010 vs</i>	<i>Female</i>	<i>13yrs vs</i>	<i>15yrs vs</i>
				<i>2002</i>	<i>2002</i>	<i>2006</i>	<i>vs Male</i>	<i>11yrs</i>	<i>11yrs</i>
<i>Austria</i>	88.0	88.0	85.9	0.834**	0.998	0.836**	0.650***	0.636***	0.564***
<i>Belgium</i>	87.7	90.3	90.1	1.265**	1.301***	0.972	0.784***	0.745***	0.606***
<i>(Flemish)</i>									
<i>Canada</i>	86.3	85.6	83.7	0.807***	0.944	0.855**	0.644***	0.945	0.942
<i>Croatia</i>	81.2	80.5	85.3	1.371***	0.963	1.424***	0.751***	0.730***	0.508***
<i>Czech Republic</i>	83.3	81.2	84.9	1.121	0.864*	1.298***	0.724***	0.883	0.883
<i>Denmark</i>	87.6	90.3	85.9	0.862*	1.343***	0.641***	0.560***	0.811**	0.856*
<i>England</i>	83.4	85.2	85.6	1.180*	1.154*	1.022	0.626***	0.867*	0.762***
<i>Estonia</i>	76.6	85.9	87.3	2.114***	1.877***	1.126	0.878***	0.743***	0.542***
<i>Finland</i>	91.6	91.5	89.9	0.813**	0.989	0.822**	0.690***	0.757***	0.610***
<i>France</i>	85.0	84.1	85.6	1.046	0.928	1.128*	0.706***	0.833***	0.669***
<i>Germany</i>	82.8	82.0	84.1	1.099	0.949	1.159**	0.711***	0.698***	0.704***
<i>Greenland</i>	91.4	82.2	84.2	0.563***	0.490***	1.149	0.596***	0.710*	0.535***
<i>Hungary</i>	84.5	81.2	82.7	0.882	0.800**	1.102	0.842***	0.927	0.590***
<i>Ireland</i>	86.4	87.4	86.6	1.033	1.106	0.934	0.714***	0.750***	0.474***
<i>Italy</i>	85.3	84.5	85.7	1.042	0.943	1.105	0.754***	0.924	0.617***

Latvia	76.9	79.1	84.6	1.662***	1.145*	1.452***	0.896*	0.909	0.802**
Lithuania	75.1	78.7	81.1	1.440***	1.225***	1.176**	0.794***	0.856**	0.715***
Macedonia	90.5	88.4	86.6	0.692***	0.798**	0.867	0.965	0.631***	0.473***
Netherlands	94.2	93.0	94.0	0.962	0.815*	1.181***	0.491***	0.727**	0.505***
Norway	82.8	88.0	87.9	1.524***	1.534***	0.993	0.669***	0.814***	0.586***
Poland	80.0	82.0	79.8	0.993	1.162**	0.855**	0.744***	0.686***	0.498***
Portugal	80.4	82.3	84.6	1.344***	1.123	1.196**	0.756***	0.776***	0.592***
Russia	76.1	79.5	82.3	1.474***	1.215***	1.213***	0.791***	0.885*	0.824***
Scotland	85.9	84.4	87.5	1.157*	0.898	1.288***	0.601***	0.729***	0.619***
Slovenia	85.6	85.7	86.8	1.090	0.998	1.092	0.677***	0.661***	0.570***
Spain	87.8	91.0	89.6	1.211**	1.446***	0.837*	0.732***	0.594***	0.459***
Sweden	85.9	87.4	86.8	1.078	1.140*	0.946	0.587***	0.547***	0.394***
Switzerland	89.1	88.2	87.4	0.846**	0.924	0.916	0.613***	0.722***	0.699***
Ukraine	74.6	81.0	79.2	1.301***	1.453***	0.895	0.910	0.962	0.661***
USA	83.0	84.1	84.2	1.077	1.075	1.002	0.760***	0.850*	0.793***
Wales	83.7	81.1	82.5	0.919	0.827**	1.110	0.556***	0.716***	0.645***

[§] The reference population was the HBSC 2010 total population

* $p < 0.05$ ** $p < 0.01$ *** $p < 0.001$